

Impaired endogenous fibrinolysis in ST-segment elevation myocardial infarction patients undergoing primary percutaneous coronary intervention is a predictor of recurrent cardiovascular events: the RISK PPCI study

Mohamed Farag^{1,2}, Nikolaos Spinthakis^{1,2}, Ying X. Gue^{1,2}, Manivannan Srinivasan², Keith Sullivan¹, David Wellsted¹, and Diana A. Gorog^{1,2,3*}

¹Postgraduate Medical School, University of Hertfordshire, Hatfield, Hertfordshire, AL10 9AB, UK; ²Department of Cardiology, East and North Hertfordshire NHS Trust, Coreys Mill Lane, Stevenage, SG1 4AB, UK; and ³Department of Medicine, National Heart & Lung Institute, Imperial College, Dovehouse Street, London SW3 6LR, UK

Received 24 March 2018; revised 29 May 2018; editorial decision 24 September 2018; accepted 15 October 2018; online publish-ahead-of-print 30 October 2018

See page 306 for the editorial comment on this article (doi: 10.1093/eurheartj/ehy760)

Aims

The endogenous fibrinolytic system serves to prevent lasting thrombotic occlusion and infarction following initiation of coronary thrombosis. We aimed to determine whether impaired endogenous fibrinolysis can identify patients with ST-segment elevation myocardial infarction (STEMI) who remain at high cardiovascular risk despite dual antiplatelet therapy (DAPT).

Methods and results

A prospective, observational study was conducted in 496 patients presenting with STEMI for primary percutaneous coronary intervention (PPCI). Blood was tested on arrival pre-PPCI, at discharge and at 30 days to assess thrombotic status using the automated point-of-care global thrombosis test and patients followed for 1 year for major adverse cardiovascular events (MACEs). Endogenous fibrinolysis was significantly impaired [baseline lysis time (LT) ≥ 2500 s] in 14% of patients and was highly predictive of recurrent MACE [hazard ratio (HR) 9.1, 95% confidence interval (CI) 5.29–15.75; $P < 0.001$], driven by cardiovascular death (HR 18.5, 95% CI 7.69–44.31; $P < 0.001$) and myocardial infarction (HR 6.2, 95% CI 2.64–14.73; $P < 0.001$), particularly within 30 days. Fibrinolysis remained strongly predictive of MACE after adjustment for conventional risk factors (HR 8.03, 95% CI 4.28–15.03; $P < 0.001$). Net reclassification showed that adding impaired fibrinolysis improved the prediction of recurrent MACE by $>50\%$ ($P < 0.001$). Patients with spontaneous ST-segment resolution pre-PPCI had more rapid, effective fibrinolysis [LT 1050 (1004–1125) s vs. 1501 (1239–1997) s, $P < 0.001$] than those without. Lysis time was not altered by standard of care STEMI treatment including DAPT and was unchanged at 30 days.

Conclusion

Endogenous fibrinolysis assessment can identify patients with STEMI who remain at very high cardiovascular risk despite PPCI and DAPT. Further studies are needed to assess whether these patients may benefit from additional, personalized antithrombotic/anticoagulant medication to reduce future cardiovascular risk.

Clinical trial registration

<http://www.clinicaltrials.gov>. Unique identifier: NCT02562690.

Keywords

STEMI • Fibrinolysis • Thrombosis • PPCI • Risk

* Corresponding author. Tel: +44 (0)207 034 8934, Fax: +44 (0)207 034 8935, Email: d.gorog@imperial.ac.uk

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2018. For permissions, please email: journals.permissions@oup.com.

Introduction

Despite primary percutaneous coronary intervention (PPCI) and optimal medications including dual antiplatelet therapy (DAPT), some 15% of patients presenting with an ST-segment elevation myocardial infarction (STEMI) experience a major adverse cardiovascular event (MACE) after the index episode, due predominantly to thrombotic complications.¹ More potent antithrombotic medications, which may reduce thrombosis, significantly increase bleeding.^{2,3} Identification of individuals at risk of future arterial thrombosis would be highly desirable, since this group could be targeted for pharmacological intervention.

The likelihood of a thrombotic event is determined by the balance between prothrombotic drivers on the one hand, and the effectiveness of the fibrinolytic system, which protects against persisting occlusive thrombosis.^{4,5} There is increasing evidence that impaired fibrinolytic capacity contributes to ongoing thrombosis risk in patients with cardiovascular disease.^{6,7} Current DAPT strategy fails to modify the residual risk of thrombosis resulting from impaired endogenous fibrinolysis. This is supported by the ATLAS-ACS-2³ and COMPASS⁸ studies showing additional risk reduction with addition of an antithrombotic/anticoagulant strategy in addition to antiplatelet therapy. Laboratory studies have shown abnormally dense fibrin structures,^{9,10} with clots that are more dense, less permeable, and more resistant to lysis in patients with stent thrombosis,¹¹ family history of premature coronary disease,¹² diabetes,^{11,13} renal failure,¹¹ and ischaemic stroke.^{11,14} However, fibrin clot structure assessment is cumbersome and lacks standardization.¹⁵ Individual components of the fibrinolytic pathway (such as t-PA, PAI-1, thrombin activatable fibrinolysis inhibitor, d-dimer) are, at best, weakly predictive of cardiovascular risk, and technical limitations and difficulty in interpretation of the relative contribution of any individual marker to overall fibrinolytic status greatly limit usefulness.¹⁶ Low plasma fibrinolytic potential, found in 10% of the population, was found to increase the risk of arterial thrombosis two-fold.¹⁷ Early data indicate that assessment of global fibrinolytic status may identify patients at increased risk of arterial thrombosis. Impaired endogenous fibrinolysis has been shown to predict MACE in non-ST elevation myocardial infarction¹⁸ and end-stage renal disease.¹⁹

The ability to predict future thrombosis would be most important in those at highest risk, such as those with STEMI, where there is a genuine need for 'personalized' treatment to reduce thrombotic complications, whilst avoiding unnecessary bleeding in low-risk groups. Blocking the thrombin pathway in addition to platelet inhibition may translate into a survival advantage, by inhibiting coagulation and improving fibrinolytic potential where this is impaired.²⁰ Our preceding pilot study in 80 STEMI patients indicated that endogenous fibrinolysis, when impaired, was associated with increased cardiovascular risk, and when enhanced, with spontaneous reperfusion.²¹ Impaired endogenous fibrinolytic status may thus identify at-risk patients, who may benefit from additional antithrombotic medication.^{6,22} We aimed to assess the usefulness of impaired endogenous fibrinolysis as a predictor of increased thrombosis risk, and as a potential target for pharmacological modulation to improve outcomes.

Methods

Study design and population

We conducted a prospective, observational, single-centre study in 496 patients in accordance with the Declaration of Helsinki and Good Clinical Practice. The study was approved by National Research Ethics Service and the UK Health Research Authority (ClinicalTrials.gov identifier: NCT02562690).

Consecutive eligible patients presenting with STEMI to our Heart Attack Centre (HAC) with a view to PPCI, were recruited. We enrolled adults (≥ 18 years) with a presumed diagnosis of STEMI based on clinical presentation and ECG criteria.²³ Patients receiving oral anticoagulation, those with known coagulation disorders, sepsis, platelet count $<100 \times 10^9/L$, haemoglobin <80 g/L, active malignancy, inability to take DAPT, or those previously enrolled in the study were excluded.

A delayed consent strategy was used, with ethical approval. In addition to routine blood tests upon arrival, an extra blood sample was taken to assess baseline thrombotic status through the same blood draw. Patients received standard-of-care antiplatelet therapy and underwent emergency angiography and PPCI as clinically indicated. Surviving patients were subsequently approached for consent. Patients who died before consent could be obtained were excluded ($n = 11$).

Antiplatelet therapy consisted of aspirin 300 mg and either clopidogrel 600 mg or ticagrelor 180 mg orally in the ambulance or emergency department upon diagnosis. Patients receiving clopidogrel pre-arrival received additional ticagrelor 180 mg loading peri-PPCI and continued on this post-procedure. Unfractionated heparin 70–100 IU/kg was given immediately before PPCI. Use of the glycoprotein IIb/IIIa inhibitor (GPI) tirofiban (Aggrastat, Corveio UK Ltd, London, UK) or bivalirudin (Angiomax, The Medicines Company, Parsippany, NJ, USA), decisions regarding access site, thrombus aspiration, and stent type were left to physician preference.

Blood sampling

Non-fasting blood samples were taken at three time-points: (i) baseline upon arrival, after DAPT loading, prior to heparin, or GPI/bivalirudin administration and before PPCI, (ii) at clinical stabilization, just prior to hospital discharge, and (iii) at 30-days' follow-up. The rationale for measuring thrombotic status at discharge and 30 days was to assess for any change over time that may be explained by an acute phase response or modification of thrombotic status by DAPT. Initial samples were taken from a 6-Fr radial or femoral sheath, which was flushed with non-heparinized saline before insertion. Subsequent samples were taken from an antecubital vein using an 18-G butterfly cannula. A two-syringe technique was employed, using the first 5 mL for routine tests and the second 5 mL for thrombotic status assessment, avoiding prolonged tourniquet time. Prior data showed no difference in thrombotic status between simultaneously collected arterial and venous samples.

Assessment of thrombotic status

Global thrombosis test

The global thrombosis test (GTT) (Thromboquest Ltd, London, UK) assesses both platelet reactivity and endogenous fibrinolysis from native, non-anticoagulated blood, and the principle of the technique has previously been described.¹⁸ The instrument was positioned in the catheterization laboratory, ready to use. The blood sample was introduced into the GTT cartridge within 15 s of withdrawal and the automated measurement begun. The instrument assesses the time taken to form an occlusive thrombus under high-shear stress (occlusion time, OT), and the time required to restore flow through endogenous fibrinolysis (lysis time, LT). The inter- and intra-assay coefficients of variation

(CV) were assessed by testing 10 stable patients on two occasions, 48 h apart and with samples run on two simultaneous channels of the instrument.

Thromboelastography

In a subgroup of 60 patients, to assess the relationship between clot strength in the thromboelastography (TEG) and lysis in the GTT, blood was also tested with thromboelastography (TEG 5000 Hemostasis Analyser System, Haemonetics, Watford, UK). Two TEG tests were performed in parallel for each patient; whole blood and whole blood plus kaolin.

ECG and angiographic analyses

ECG and angiographic analyses were performed by an investigator blinded to thrombotic status results and not involved in the PPCI procedure. Standard 12-lead ECGs obtained in the ambulance or emergency department (baseline), and on arrival to the HAC were analysed using a hand-held caliper. The ST-segment was measured 20 ms after the J-point, and the sum of ST deviation measured as previously described.²⁴ The percent resolution of ST deviation from baseline to arrival was categorized using Schröder's three-component definition: complete ($\geq 70\%$), partial (30–70%), and no ($\leq 30\%$) ST-segment resolution. Flow in the infarct-related artery was graded using the Thrombolysis in Myocardial Infarction (TIMI) classification and patency defined as TIMI 2 or 3 flow.

Scanning electron microscopy

In a cohort of representative patients, two blood samples per patient ($n = 10$) at the time of acute STEMI presentation were run in parallel channels in the GTT. Whilst in the first channel the measurement was allowed to proceed as normal, in the second channel, the test was terminated after OT. The thrombus formed between the two ball-bearings was carefully extracted, immediately washed in Na-cacodylate buffer and fixed with 2.5% glutaraldehyde, before point-critical dehydration with ethanol (5–100%) and hexamethyldisilazane. Samples were coated with gold palladium and imaged using a Phenom ProX (EDS) scanning electron microscope (SEM) (Lambda Photometrics Ltd, Hertfordshire, UK).

Data collection and follow-up

During the index admission, case notes and electronic records were examined, to allow contemporaneous completion of study-specific case record forms. Patients were followed-up at 30-days in person, and at 6 and 12 months by telephone and by accessing casenotes.

Study endpoints

The primary endpoint was the occurrence of MACE, defined as the composite of cardiovascular death (CVD), nonfatal myocardial infarction (MI) including stent thrombosis (defined according to the Academic Research Consortium criteria), or stroke. These are defined in the [Supplementary material online](#). The secondary endpoint was major bleeding classified as type 3–5 according to the Bleeding Academic Research Consortium (BARC) definition. For all endpoints, source documents were obtained and diagnosis verified by two independent clinicians blinded to thrombotic status results.

Statistical analysis

This study aimed to assess whether LT is a predictor of MACE in patients with STEMI. Based on an effect size (hazard ratio, HR) of 3.3 from our pilot,²¹ assuming a two-sided alpha of 0.01, 10% MACE and 10% attrition rate, using the Cox proportional hazard model we calculated that 464

patients would be required to achieve 90% power. A lower α level and higher power were employed, to adequately address the study question.

Data are presented as mean (standard deviation) or median and inter-quartile range (IQR). Dichotomous variables were compared using χ^2 test or Fisher's exact test. Correlations were analysed using Spearman's method. Ability of the test to discriminate between patients with and without the study endpoint MACE (a composite of CVD, nonfatal MI, or stroke) was evaluated by receiver-operating characteristic (ROC) curve analysis. The Kaplan–Meier survival methods with log-rank tests were used. Only events occurring beyond the point of LT testing were related to the test's ability to predict MACE. To investigate the relationship between LT and MACE, univariate and multivariable hazard regression models of Cox were used. The bootstrap technique using one thousand samples was used as a way to account for final multivariable model uncertainty. All study variables were first analysed with univariate analysis and those that showed a significant interaction ($P < 0.05$) were entered into the multivariable analysis. Multiple fractional polynomial models were used to illustrate the nonlinear association between LT and MACE. Net reclassification improvement (NRI) was used to assess the change in discrimination after adding LT ≥ 2500 to the combination of the three baseline predictors (age, diabetes, and TIMI score). Analyses were performed with Stata V.15 (StataCorp, College Station, TX, USA).

Results

Between April 2015 and March 2017, 543 patients were eligible for recruitment. Six patients were already enrolled in the trial and 41 met ≥ 1 exclusion criteria. All survivors were followed for a median of 12 months. Clinical characteristics are shown in [Table 1](#) and baseline bloodwork, angiographic, procedural, and echocardiographic characteristics in [Supplementary material online, Tables S2 and S3](#).

Thrombotic status over 30-days

The inter- and intra-assay CVs were 8% and 9% for OT and 10% and 10% for LT, respectively. There was a negative correlation between baseline OT and baseline LT ($r = -0.4$, $P < 0.001$). Compared with baseline, OT increased significantly at hospital discharge and 30-days ([Supplementary material online, Figure S1A](#)). Compared with baseline, LT did not change at discharge ($P = 0.761$) or 30-days ($P = 0.652$). Baseline LT correlated with LT at discharge ($r = 0.2$, $P < 0.001$) and LT at 30-days ($r = 0.2$, $P < 0.001$) ([Supplementary material online, Figure S1B](#)).

Relationship between patient characteristics and thrombotic status

Patient characteristics in [Table 1](#) and [Supplementary material online, Tables S1 and S2](#) were interrogated for relation to baseline OT and LT. Baseline OT was related to pre-PPCI TIMI-3 flow ($r = 0.2$, $P < 0.001$), peak troponin ($r = -0.2$, $P = 0.002$), and neutrophil count ($r = -0.2$, $P < 0.001$). Baseline LT was related to pre-PPCI TIMI-3 flow ($r = -0.2$, $P < 0.001$). Impaired endogenous fibrinolysis (baseline LT ≥ 2500 s) was significantly more frequent in patients with cardiogenic shock, prior MI or percutaneous coronary intervention (PCI), low albumin, raised fibrinogen, raised high-sensitivity C-reactive protein, and high TIMI score ([Table 1, Supplementary material online, Tables S1 and S2](#)). There was a very weak but significant correlation between baseline LT and peak troponin ($r = 0.14$, $P = 0.016$). There was

Table 1 Baseline patient characteristics

	Whole group (n = 496)	Baseline LT <2500 (n = 426)	Baseline LT ≥2500 (n = 70)	P-value
Age (years)	63 (55–74)	63 (54–72)	67 (56–79)	0.082
Male	386 (77.8)	332 (77.9)	54 (77.1)	0.877
Caucasian	447 (90.1)	382 (89.7)	65 (92.9)	0.710
BMI	26.7 (23.9–30.1)	26.8 (23.9–30.2)	26.1 (23.7–29.9)	0.487
Cardiogenic shock ^a	18 (3.6)	12 (2.8)	6 (8.6)	0.029
TIMI score	3 (1–5)	3 (1–4)	4 (1–6)	<0.001
GRACE score	115 (92–140)	115 (92–135)	125 (101–174)	0.011
Diabetes mellitus	96 (19.4)	77 (18.1)	19 (27.1)	0.101
Active smoker	164 (33.1)	137 (32.2)	27 (38.6)	0.337
Hypertension	247 (49.8)	213 (50.0)	34 (43.6)	0.898
Family history of premature IHD	206 (41.5)	178 (41.8)	28 (40.0)	0.795
Prior MI	61 (12.3)	47 (11.0)	14 (20.0)	0.048
Prior PCI	56 (11.3)	41 (9.6)	15 (21.4)	0.007
Prior CABG	7 (1.4)	5 (1.2)	2 (2.9)	0.258
Renal insufficiency	25 (5.0)	21 (4.9)	4 (5.7)	0.768
PVD	20 (4.0)	17 (3.9)	3 (4.3)	0.753
Prior CVA	23 (4.6)	17 (3.9)	6 (8.6)	0.117
Prior statin use	134 (27.0)	114 (26.8)	20 (28.6)	0.772
Prior aspirin use	91 (18.3)	73 (17.1)	18 (25.7)	0.096
Prior P2Y ₁₂ inhibitor use	22 (4.4)	16 (3.8)	6 (8.6)	0.069
Initial P2Y ₁₂ inhibitor loading agent				
Clopidogrel	407 (82.1)	347 (81.5)	59 (84.3)	0.620
Ticagrelor	78 (15.7)	69 (16.2)	9 (12.9)	0.491
Cangrelor	11 (2.2)	9 (2.1)	2 (2.9)	1.000
Morphine prior to blood sample	327 (65.9)	276 (64.8)	51 (72.9)	0.221
Time from P2Y ₁₂ inhibitor loading to first blood sample (min)	46 (37–60)	45 (36–60)	50 (40–60)	0.145
Medications prior to hospital discharge				
Aspirin	472 (95.2)	408 (95.8)	64 (91.4)	0.130
Clopidogrel	81 (16.3)	68 (15.9)	13 (18.6)	0.601
Ticagrelor	388 (78.2)	337 (79.1)	51 (72.9)	0.274
Prasugrel	2 (0.4)	2 (0.5)	0	1.000
Beta-blocker	438 (88.3)	380 (89.2)	58 (82.9)	0.157
ACE inhibitor	433 (87.3)	376 (88.3)	57 (81.4)	0.123
Calcium antagonist	29 (5.8)	26 (6.1)	3 (4.3)	0.606
Statin	466 (93.9)	403 (94.6)	63 (90.0)	0.169
Nitrate	18 (3.6)	15 (3.5)	3 (4.3)	0.999
Insulin	42 (8.5)	35 (8.2)	7 (10.0)	0.642

Values are presented as median (IQR) or n (%). Renal insufficiency defined as creatinine >177 µmol/L. Prior statin, aspirin, or P2Y₁₂ inhibitor defined as regular use pre-hospitalization. Family history of premature IHD defined as a diagnosis of IHD in a first-degree relative <60 years. Values in bold are significant (i.e. $P < 0.05$).

ACE, angiotensin-converting enzyme; BMI, body mass index; CABG, coronary artery bypass grafting; CVA, cerebrovascular accident; IHD, ischaemic heart disease; LT, lysis time; MI, myocardial infarction; PCI, percutaneous coronary intervention; PVD, peripheral vascular disease; TIMI, Thrombolysis in Myocardial Infarction.

^aPatients artificially ventilated or with reduced conscious level received rectal aspirin 300 mg and either intravenous cangrelor 30 mcg/kg bolus followed by 4 mcg/kg/min infusion ($n = 11$), or ticagrelor 180 mg via nasogastric tube ($n = 7$) on arrival.

no correlation between baseline OT or baseline LT and time of day of blood sampling or pain-to-balloon time.

Thrombotic status and cardiovascular outcomes

There were 53 MACE events among 46 patients (Table 2). Baseline OT was inversely related to MACE with a c-statistic of

0.32 [95% confidence interval (CI) 0.233–0.407] with a cut-point of 302 s, with sensitivity 71% and specificity 67.5% (Figure 1A). Baseline OT < 302 s was related to MACE (HR 0.34, 95% CI 0.191–0.594; $P = 0.0001$). Patients who experienced MACE demonstrated baseline OT that was significantly shorter (more thrombotic) than patients without MACE [268 (180–333) s vs. 367 (278–484) s; $P < 0.001$], particularly with respect to MI [231 (106–300) s vs. 363 (275–483) s; $P < 0.001$], and ischaemic stroke [186

Table 2 Clinical outcomes at 12 months follow-up

	Whole group (n = 496)	Baseline LT <2500 (n = 426)	Baseline LT ≥2500 (n = 70)	HR (95% CI)	P-value
MACE (cardiovascular death, re-infarction, CVA)	53 (10.7)	23 (5.4)	30 (42.9)	9.1 (5.29–15.75)	<0.001
Cardiovascular death	25 (5.0)	7 (1.6)	18 (25.7)	18.5 (7.69–44.31)	<0.001
New myocardial infarction or re-infarction	21 (4.2)	11 (2.8)	10 (14.3)	6.23 (2.64–14.73)	<0.001
Definite stent thrombosis					
Acute (<24 h)	6 (1.2)	3 (0.7)	3 (4.3)	5.58 (1.13–27.63)	0.046
Subacute (24 h to 30 days)	4 (0.8)	2 (0.5)	2 (2.9)	7.32 (1.02–52.14)	0.064
Late (>30 days)	3 (0.6)	2 (0.5)	1 (1.4)	4.36 (0.39–48.34)	0.281
Cerebrovascular accident	7 (1.4)	5 (1.2)	2 (2.9)	2.71 (0.52–14.01)	0.273
Major bleeding (BARC Type 3–5)	7 (1.4)	5 (1.2)	2 (2.9)	3.23 (0.62–16.70)	0.204
Minor bleeding (BARC Type ≤2)	19 (3.8)	17 (3.9)	2 (2.9)	1.03 (0.24–4.47)	0.967
Non cardiovascular death	9 (1.8)	7 (1.6)	2 (2.9)	2.42 (0.50–11.68)	0.312

Values are presented as n (%). Values in bold are significant (i.e. $P < 0.05$).

BARC, Bleeding Academic Research Consortium; CI, confidence interval; CVA, cerebrovascular accident; HR, hazard ratio; LT, lysis time; MACE, major adverse cardiovascular events.





Figure 1 Receiver-operating characteristic curves for occlusion time (A) and lysis time (B) for recurrent adverse events to 365 days. Baseline lysis time ≥ 2500 s predicted major adverse cardiovascular events, with 60% sensitivity and 91% specificity. For occlusion time at baseline risk increases with falling occlusion time. To provide the standard monotonically increasing receiver-operating characteristic curve function coding for events was therefore reversed.

(125–288) s vs. 360 (270–482) s; $P = 0.003$] (Supplementary material online, Figure S2). The OT at hospital discharge did not differ between patients who experienced MACE or major bleed and those who did not (Figure 1A).


Baseline LT significantly discriminated between patients with and without MACE, with a c -statistic of 0.78 (95% CI 0.691–0.859; $P < 0.05$) (Figure 1B). Lysis time ≥ 2483 s was the optimal cut-point to predict MACE (rounded to 2500 s for ease), with sensitivity 60% and



specificity 90.5% (Figure 1B). There were no LT readings between 2483 s and 2500 s that could have been wrongly classified based on rounding the cut-point. Baseline LT ≥ 2500 s was strongly related to MACE (HR 9.1, 95% CI 5.29–15.75; $P < 0.001$) (Figure 2), driven by CVD (HR 18.5, 95% CI 7.69–44.31; $P < 0.001$) and MI (HR 6.2, 95% CI 2.64–14.73; $P < 0.001$) (Figures 3B and 4) (Table 2) and probability of MACE increased with increasing baseline LT (Figures 2B and 5). Lysis time at subsequent timepoints was not related to MACE.

Prognostic value of impaired endogenous lysis time

Of the patient characteristics (Table 1, Supplementary material online, Tables S1 and S2), the following were related to MACE: age ($P < 0.001$), prior MI ($P = 0.003$), prior PCI ($P = 0.013$), prior stroke ($P = 0.016$), prior statin (0.005), prior aspirin ($P = 0.002$), cardiogenic shock ($P = 0.003$), TIMI score ($P < 0.001$), low haemoglobin ($P < 0.001$), low haematocrit ($P = 0.001$), low albumin ($P < 0.001$), high creatinine ($P = 0.009$), high C-reactive protein ($P = 0.003$), short



baseline OT ($P < 0.001$), number of diseased vessels on angiography ($P = 0.007$), left ventricular dysfunction ($P < 0.001$), and absence of spontaneous ST-resolution pre-PPCI ($P = 0.032$) (Supplementary material online, Table S4).

The following were then entered into the final multivariable Cox proportional hazard model: age, prior stroke, prior statin, prior aspirin, creatinine, number of diseased vessels, and left ventricular dysfunction. None of these covariates were significantly correlated either with LT or its dichotomized version. Although low haemoglobin and haematocrit levels were not related to LT, these were not included in the model, as we felt they may be related at the extremes. Baseline LT ≥ 2500 s remained significantly predictive of MACE after adjustment for other risk factors (HR 8.03, 95% CI 4.28–15.03; $P < 0.001$) and after bootstrap resampling, the optimism-corrected c-index was -0.015 and the calibration slope was 0.933 which suggests no over fitting of the model. It remained significant after the addition of diabetes into the final model (HR 8.44, 95% CI 4.47–15.93; $P < 0.001$) and also after excluding patients with cardiogenic




Figure 4 Multiple fractional polynomial curves showing major adverse cardiovascular event rates according to baseline lysis time. Solid lines are shown as estimate and 95% confidence interval. Dotted line indicates lysis time ≥ 2500 s.

shock (HR 8.07, 95% CI 4.55–14.34; $P < 0.001$) or severe LV dysfunction (HR 7.14, 95% CI 3.77–13.49; $P < 0.001$) from the analysis.

Net reclassification improvement showed that inclusion of baseline LT ≥ 2500 s in a model containing three baseline predictors (age, diabetes, and TIMI score) significantly added to the model effectiveness (NRI estimate 0.514, $P < 0.001$).

Spontaneous coronary reperfusion and ST-resolution

Patients with spontaneous reperfusion (Supplementary material online, Table S2), manifesting as complete or partial resolution of ST-elevation pre-PPCI, all had TIMI 2–3 flow at presentation, shorter baseline LT [1050 (1004–1125) s vs. 1501 (1239–1997) s, $P < 0.001$] and longer baseline OT [448 (371–586) s vs. 338 (245–454) s, $P < 0.001$] than patients without spontaneous reperfusion. Spontaneous reperfusion correlated with OT ($r = 0.3$, $P < 0.001$) and inversely with LT ($r = -0.4$, $P < 0.001$).

Scanning electron microscopy

Increasing LT *in vitro* was associated with increasing density of the fibrin network on SEM (Figure 6) and the association was discernible in each tested patient. Images are illustrative of the possible pathophysiological mechanism underlying the observed resistance to lysis and should be interpreted in the context of the limitations pertaining to the small number of samples analysed as well as possible inter-observer variability.

Thromboelastography

Baseline OT correlated with the following whole blood TEG indices (explained in Supplementary material online, Table S3): R ($r = 0.4$, $P = 0.002$), K ($r = 0.2$, $P = 0.049$), TMA ($r = 0.3$, $P = 0.027$), and MA ($r = -0.2$, $P = 0.050$). Similar relationships were observed with kaolin:




Figure 5 Major adverse cardiovascular events according to baseline lysis time. Hazard ratios for cardiovascular death, myocardial infarction, and cerebrovascular accident. The 95% confidence interval shown in brackets at top of bars. * $P < 0.001$. CVA, cerebrovascular accident; CVD, cardiovascular death; E, number of events; MI, myocardial infarction; n, number of patients.

R ($r = 0.4$, $P = 0.001$), K ($r = 0.4$, $P = 0.009$), TMA ($r = 0.3$, $P = 0.038$), and MA ($r = -0.4$, $P = 0.005$). Baseline LT did not correlate with any TEG indices, including LY30 and LY60 using whole blood with or without kaolin.

Discussion

In patients with STEMI presenting for PPCI, impaired endogenous fibrinolysis detected using an automated point-of-care test in the catheterization laboratory, is a novel risk factor for subsequent MACE with significant predictive value (Take home figure).

Whilst platelet reactivity fell from admission to hospital discharge and further by day 30 (shown by increasing OT), likely representing the effects of antiplatelet medications and plaque stabilization, endogenous fibrinolysis by comparison, did not change from admission to 30-days. Endogenous fibrinolysis was significantly impaired in 14% of patients and was highly predictive of recurrent MACE, driven by CVD and MI. As in other contemporary PPCI trials, most events occurred in hospital, and the majority of the rest within the first 30-days, and thus because of the small number of events, the study was underpowered to assess the predictive power of LT measured at discharge and 30-days. Prolonged LT was associated with increasing fibrin network density with reduced pore size on SEM, which likely explains the resistance to lysis, in line with earlier studies.^{7,9,11,12}

Our data showing impaired endogenous fibrinolysis to be highly predictive of outcome, confirms findings from our STEMI pilot²¹ and is supported by data in a lower-risk cohort with non-ST elevation MI, where impaired fibrinolysis was predictive of increased cardiovascular risk (HR 2.5).¹⁸ Our findings are strongly supported by a very recent publication analysing stored plasma samples from >4000 ACS patients in the PLATO trial, showing that clot lysis time, measured




Figure 6 Composition of *in vitro* formed thrombus at the time of acute myocardial infarction. Scanning electron microscope $\times 1400$ (upper panel) and $\times 9000$ (lower panel) magnification, from patients with lysis time 990 s (A1, A2), lysis time 3160 s (B1, B2), and lysis time >6000 s (C1, C2). Increasing lysis time *in vitro* was associated with increasing density of the fibrin network.


using a turbidimetric assay, was an independent predictor of adverse outcome, with risk increasing as lysis time increased, independent of antiplatelet medications.²⁵ However, unlike the specialized turbidimetric assay performed in a centralized laboratory, we report results with a point-of-care test that could influence clinician decision making in a timely manner, allowing personalization of treatment before hospital discharge to reduce 30-day MACE.

The lack of a linear relationship between LT and outcome likely reflects the small number of patients in the 3rd and 4th LT quartiles ($n = 21$ and $n = 30$, respectively), such that the error of estimation is large and compromises the ability to evaluate the relationship between the magnitude of LT and the risk of MACE. However, we postulate that there would have been a linear relationship had we had a larger sample size.


There were no cardiovascular risk factors that clearly related to LT, but numbers were too small for subgroup analysis. However, amongst patients with very high-risk, namely those with cardiogenic shock, high TIMI scores, low albumin, and high C-reactive protein, significantly more patients exhibited impaired fibrinolysis than lower risk patients. The relationship between cardiogenic shock and

prolonged LT is supported by data from the recently published PLATO substudy also showing an association between NT-proBNP levels and lysis time.²⁵ Whilst we did not measure NT-proBNP, the proportion of patients with cardiogenic shock increased with increasing lysis time. Such patients are known to be at very high risk. However, LT remained a significant predictor of recurrent MACE even after excluding patients with cardiogenic shock. Unlike the PLATO substudy, we saw only a weak relationship between troponin and LT. However, any conclusions drawn from this have to be tempered by the unusual observation that peak troponin was not predictive of outcome in our population. Impaired fibrinolysis appears not to simply mirror conventional cardiovascular risk profile, but to be a novel risk factor, which when added to traditional risk factors very significantly improved the prediction of MACE.

An important issue is whether impaired endogenous fibrinolysis is a modifiable risk factor. Whilst antiplatelet medications reduce platelet reactivity over time, LT appears unaffected by oral antiplatelet medications, or plaque stabilization. Previous work suggests that non-vitamin K antagonist oral anticoagulants (NOACs)²⁶ and the PAR-1 inhibitor vorapaxar²⁷ can favourably modulate (reduce) LT.



* Impaired endogenous fibrinolysis associated with significantly increased risk of MACE especially in first 30 days post-STEMI, driven by cardiovascular death and re-infarction



Take home figure Impaired endogenous fibrinolysis in patients with ST-segment elevation myocardial infarction presenting for primary percutaneous coronary intervention (PPCI), detected using an automated point-of-care technique in the catheterization laboratory, identifies those at high risk of recurrent major adverse cardiovascular events despite dual antiplatelet therapy and PPCI.

The COMPASS trial showed that among patients with stable atherosclerosis, ultra-low dose rivaroxaban plus aspirin reduced the risk of the composite of CVD, stroke, or MI and more major bleeding compared with aspirin alone.⁸ In ACS patients, addition of a NOAC, such as rivaroxaban in ATLAS-ACS-2³ or vorapaxar² to DAPT can reduce the risk of recurrent ischaemic events, although offset by a significant increase in bleeding. Since this strategy clearly cannot be recommended for all ACS patients, assessing endogenous fibrinolysis could identify those at highest risk, who may benefit most from additional anticoagulation.

Spontaneous reperfusion was observed in 15% of patients pre-PPCI, who had more rapid (efficient) endogenous fibrinolysis and reduced platelet reactivity, supporting the signal seen in our pilot. We postulate that this may indicate a cohort of patients where complete thrombotic vessel occlusion can occur due to a very potent thrombotic stimulus, but in whom endogenous fibrinolysis is more effective than in others, sufficient to spontaneously restore coronary flow.

In our sub-study, clot lysis with TEG did not relate to fibrinolysis with the GTT. Thromboelastography is a global test of coagulation, simultaneously assessing clot development, stabilization and dissolution and is useful for assessing clot strength, but its value in assessing spontaneous fibrinolytic status is questionable, as it fails to assess the procoagulant (thrombin-generating) and fibrinolysis-inhibiting (PAI-1, TAFI) properties of platelets.⁶ It more closely resembles the low-shear environment of venous stasis than the high-shear situation in the GTT, which likely explains the lack of correlation between these tests.

Although baseline fibrinogen levels were higher in patients with impaired fibrinolysis, fibrinogen is an acute phase protein and levels

normalize following an acute event. C-reactive protein and albumin may similarly reflect acute illness. On the other hand, endogenous fibrinolysis did not change over time. Previous studies have measured individual proteins involved in fibrinolysis, including t-PA, PAI-1, alpha-2 antiplasmin, alpha2-antiplasmin-plasmin complex, D-dimer, TAFI, and Lp(a). Technical limitations, combined with challenges in knowing the relative importance and contribution of individual variables to overall fibrinolytic status, limit the usefulness of this approach.¹⁶ In this study, the predictive power of endogenous fibrinolytic activity was considerably higher than that of plasma PAI-1,^{28,29} perhaps because PAI-1 release from the platelet core of a thrombus is localized, and not necessarily reflected by systemic PAI-1 antigen levels.

Limitations

The main limitations of our study are the sample size and recruitment from a single centre. The numbers are too small to assess relationships between endogenous fibrinolysis and established cardiovascular risk factors. Because of the need to obtain informed consent, some of the sickest patients who died before giving consent, were excluded. Optimal antithrombotic medication for STEMI comprises of aspirin with ticagrelor or prasugrel, but due to financial constraints in much of the UK, the ambulance crew often administer clopidogrel and aspirin. However, such patients were additionally given ticagrelor on arrival and it is unlikely that prasugrel or ticagrelor administration in the ambulance instead of clopidogrel would have impacted on endogenous fibrinolysis. Baseline samples were obtained before the

acutely administered antiplatelet agents would have had maximal effect and so reflect real-time thrombotic status at the time of PPCI. The study was underpowered to assess the relationship between thrombotic status and bleeding events, since very few bleeds were observed at least in part due to the majority of interventions being performed radially and the exclusion of anticoagulated patients.

Conclusion

Impaired endogenous fibrinolysis in patients with STEMI presenting for PPCI, detected using an automated point-of-care technique in the catheterization laboratory, identifies those at high risk of recurrent MACE. This novel risk factor improved the prediction of recurrent MACE by >50% when added to conventional predictors. Further studies are needed to assess whether impaired endogenous fibrinolysis is a modifiable risk factor, targetable by pharmacotherapy, to reduce cardiovascular risk.

Supplementary material

Supplementary material is available at *European Heart Journal* online.

Acknowledgements

We would like to gratefully acknowledge the help of Dr Gareth Ambler, Senior Lecturer in Medical Statistics at University College, London for providing us with statistical assistance.

Funding

The project was funded by the R&D department of East and North Hertfordshire NHS Trust.

Conflict of interest: D.A.G. is related through family to a company director in Thromboquest Ltd, which manufactures the Global Thrombosis Test, but neither she, nor her spouse, nor children have financial involvement or equity interest in and have received no financial assistance, support, or grants from Thromboquest Ltd. Thromboquest Ltd had no involvement in design, conduct, or finance of this study and the instrument and disposables were obtained through a normal commercial transaction. D.A.G. has received grants/honoraria from Bayer, BMS, Astra Zeneca, and Abbott. All other authors declared no conflict of interest.

References

- Giustino G, Mehran R, Dangas GD, Kirtane AJ, Redfors B, G  n  reux P, Brener SJ, Prats J, Pocock SJ, Deliangryis EN, Stone GW. Characterization of the average daily ischemic and bleeding risk after primary PCI for STEMI. *J Am Coll Cardiol* 2017;**70**:1846–1857.
- Tricoci P, Huang Z, Held C, Moliterno DJ, Armstrong PW, Van de Werf F, White HD, Aylward PE, Wallentin L, Chen E, Lokhnygina Y, Pei J, Leonardi S, Rorick TL, Kilian AM, Jennings LH, Ambrosio G, Bode C, Cequier A, Cornel JH, Diaz R, Erkan A, Huber K, Hudson MP, Jiang L, Jukema JW, Lewis BS, Lincoff AM, Montalescot G, Nicolau JC, Ogawa H, Pfisterer M, Prieto JC, Ruzyllo W, Sinnaeve PR, Storey RF, Valgimigli M, Whellan DJ, Widimsky P, Strony J, Harrington RA, Mahaffey KW. Thrombin-receptor antagonist vorapaxar in acute coronary syndromes. *N Engl J Med* 2012;**366**:20–33.
- Mega JL, Braunwald E, Wiviott SD, Bassand JP, Bhatt DL, Bode C, Burton P, Cohen M, Cook-Bruns N, Fox KA, Goto S, Murphy SA, Plotnikov AN, Schneider D, Sun X, Verheugt FW, Gibson CM. Rivaroxaban in patients with a recent acute coronary syndrome. *N Engl J Med* 2012;**366**:9–19.
- Barrab  s JA, Galian L. Endogenous thrombolysis: a hidden player in acute coronary syndromes? *J Am Coll Cardiol* 2010;**55**:2116–2117.
- Lucking AJ, Gibson KR, Paterson EE, Faratian D, Ludlam CA, Boon NA, Fox KA, Newby DE. Endogenous tissue plasminogen activator enhances fibrinolysis and limits thrombus formation in a clinical model of thrombosis. *Arterioscler Thromb Vasc Biol* 2013;**33**:1105–1111.
- Okafor O, Gorog D. Endogenous fibrinolysis: an important mediator of thrombus formation and cardiovascular risk. *J Am Coll Cardiol* 2015;**65**:1683–1699.
- Leander K, Blomb  ck M, Wall  n H, He S. Impaired fibrinolytic capacity and increased fibrin formation associate with myocardial infarction. *Thromb Haemost* 2012;**107**:1092–1099.
- Eikelboom JW, Connolly SJ, Bosch J, Dagenais GR, Hart RG, Shestakovska O, Diaz R, Alings M, Lonn EM, Anand SS, Widimsky P, Hori M, Avezum A, Piegas LS, Branch KRH, Probstfield J, Bhatt DL, Zhu J, Liang Y, Maggioni AP, Lopez-Jaramillo P, O'Donnell M, Kakkar AK, Fox KAA, Parkhomenko AN, Ertl G, St  rk S, Keltai M, Ryden L, Pogosova N, Dans AL, Lanas F, Commerford PJ, Torp-Pedersen C, Guzik TJ, Verhamme PB, Vinereanu D, Kim JH, Tonkin AM, Lewis BS, Felix C, Yusuf S, Steg PG, Mehta SR, Cook-Bruns N, Mittleman B, Chen E, Leong D, Yusuf S. Rivaroxaban with or without aspirin in stable cardiovascular disease. *N Engl J Med* 2017;**377**:1319–1330.
- Collet JP, Allali Y, Lesty C, Tanguy ML, Silvain J, Ankri A, Blanchet B, Dumaine R, Gianetti J, Payot L, Weisel JW, Montalescot G. Altered fibrin architecture is associated with hypofibrinolysis and premature coronary atherosclerosis. *Arterioscler Thromb Vasc Biol* 2006;**26**:2567–2573.
- Fatah K, Hamsten A, Blomb  ck B, Blomb  ck M. Fibrin gel network characteristics and coronary heart disease: relations to plasma fibrinogen concentration, acute phase protein, serum lipoproteins and coronary atherosclerosis. *Thromb Haemost* 1992;**68**:130–135.
- Undas A. Fibrin clot properties and their modulation in thrombotic disorders. *Thromb Haemost* 2014;**112**:32–42.
- Mills JD, Ari  ns RA, Mansfield MW, Grant PJ. Altered fibrin clot structure in the healthy relatives of patients with premature coronary artery disease. *Circulation* 2002;**106**:1938–1942.
- Alzahrani SH, Hess K, Price JF, Strachan M, Baxter PD, Cubbon R, Phoenix F, Gamlen T, Ari  ns RA, Grant PJ, Ajjan RA. Gender-specific alterations in fibrin structure function in type 2 diabetes: associations with cardiometabolic and vascular markers. *J Clin Endocrinol Metab* 2012;**97**:E2282–E2287.
- Rooth E, Wall  n NH, Blomb  ck M, He S. Decreased fibrin network permeability and impaired fibrinolysis in the acute and convalescent phase of ischemic stroke. *Thromb Res* 2011;**127**:51–56.
- Pieters M, Undas A, Marchi R, De Maat MP, Weisel J, Ari  ns RA. An international study on the standardization of fibrin clot permeability measurement: methodological considerations and implications for healthy control values. *J Thromb Haemost* 2012;**10**:2179–2181.
- Gorog DA. Prognostic value of plasma fibrinolysis activation markers in cardiovascular disease. *J Am Coll Cardiol* 2010;**55**:2701–2709.
- Guimar  es AH, de Bruijne EL, Lisman T, Dippel DW, Deckers JW, Poldermans D, Rijken DC, Leebeek FW. Hypofibrinolysis is a risk factor for arterial thrombosis at young age. *Br J Haematol* 2009;**145**:115–120.
- Saraf S, Christopoulos C, Salha IB, Stott DJ, Gorog DA. Impaired endogenous thrombolysis in acute coronary syndrome patients predicts cardiovascular death and nonfatal myocardial infarction. *J Am Coll Cardiol* 2010;**55**:2107–2115.
- Sharma S, Farrington K, Kozarski R, Christopoulos C, Niespialowska-Steuden M, Moffat D, Gorog DA. Impaired thrombolysis: a novel cardiovascular risk factor in end-stage renal disease. *Eur Heart J* 2013;**34**:354–363.
- Capodanno D. Oral antithrombotic therapy after acute coronary syndromes: “dual antiplatelet” or “dual pathway”? *EuroIntervention* 2017;**13**:773–775.
- Christopoulos C, Farag M, Sullivan K, Wellsted D, Gorog DA. Impaired thrombolytic status predicts adverse cardiac events in patients undergoing primary percutaneous coronary intervention. *Thromb Haemost* 2017;**117**:457–470.
- Gorog DA, Fayad ZA, Fuster V. Arterial thrombus stability: does it matter and can we detect it? *J Am Coll Cardiol* 2017;**70**:2036–2047.
- Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology (ESC). ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2012;**33**:2569–2619.
- de Lemos JA, Antman EM, Giugliano RP, McCabe CH, Murphy SA, Van de Werf F, Gibson CM, Braunwald E. ST-segment resolution and infarct-related artery patency and flow after thrombolytic therapy. Thrombolysis in Myocardial Infarction (TIMI) 14 investigators. *Am J Cardiol* 2000;**85**:299–304.
- Sumaya W, Wallentin L, James SK, Siegbahn A, Gabrysch K, Bertilsson M, Himmelmann A, Ajjan RA, Storey RF. Fibrin clot properties independently predict adverse clinical outcome following acute coronary syndrome: a PLATO sub-study. *Eur Heart J* 2018;**39**:1078–1085.
- Farag M, Niespialowska-Steuden M, Okafor O, Artman B, Srinivasan M, Khan A, Sullivan K, Wellsted D, Gorog DA. Relative effects of different non-vitamin K antagonist oral anticoagulants on global thrombotic status in atrial fibrillation. *Platelets* 2016;**27**:687–693.
- Rosser G, Tricoci P, Morrow D, Christopoulos C, Niespialowska-Steuden MN, Kozarski R, Kozarski R, Wilcox R, Gorog DA. PAR-1 antagonist vorapaxar

favorably improves global thrombotic status in patients with coronary disease. *J Thromb Thrombolysis* 2014;**38**:423–429.

28. Song C, Burgess S, Eicher JD, O'Donnell CJ, Johnson AD. Causal effect of plasminogen activator inhibitor type 1 on coronary heart disease. *J Am Heart Assoc* 2017;**6**:e004918.

29. Iacoviello L, Agnoli C, De Curtis A, di Castelnuovo A, Giordanella MC, Krogh V, Mattiello A, Matullo G, Sacerdote C, Tumino R, Vineis P, de Gaetano G, Panico S, Donati MB. Type 1 plasminogen activator inhibitor as a common risk factor for cancer and ischaemic vascular disease: the EPICOR study. *BMJ Open* 2013;**3**:e003725.

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehy782

Online publish-ahead-of-print 28 November 2018


Myocardial infarction after dog bite

Alberto Polimeni¹, Carmen Spaccarotella¹, Annalisa Mongiardo¹, and Ciro Indolfi^{1,2*}

¹Division of Cardiology, Department of Medical and Surgical Sciences, Magna Graecia University, Catanzaro 88100, Italy; and ²URT-CNR, Magna Graecia University, Catanzaro, Italy

* Corresponding author. Tel: +39 096 1364 7151, Fax: +39 096 1364 7153, Email: indolfi@unicz.it

An 80-year-old man with a family history of cardiovascular disease and arterial hypertension presented typical chest pain after multiple shepherd dog bites resulting in severe body injury (Panels A–C) without a concomitant infection. The electrocardiogram revealed ST-segment elevation in leads V1–V5 (Panel D) with partial resolution of ST segment after thrombolytic therapy (Panel E). The patient was transferred to our institution for urgent coronary angiography that revealed a significant stenosis of left anterior descending (LAD) artery with a thrombotic formation (Panel F) that was successfully treated with drug eluting stent implantation.



The association between myocardial infarction and dog bite is rare and described as a complication of *Capnocytophaga canimorsus* infection causing fulminant sepsis with a direct myocardial injury or endocarditis with septic emboli. However, in the case presented, the patient was not infected, as demonstrated by specific blood cultures, and no clinical signs of infection were present. Since coronary angiography revealed a significant stenosis of LAD artery with a thrombotic formation, it is reasonable to believe that the stress associated with multiple dog bites could be responsible for the transition from chronic to acute coronary syndrome. The acute rupture of a vulnerable plaque could be the consequence of an abrupt sympathetic stimulation. Increase in ventricular inotropy, heart rate, blood pressure, or stimulation of alpha-adrenergic receptors may alter the shear stress, contributing to the rupture of the shoulder of vulnerable atherosclerotic plaque.

To date, this is the first report, to the best of our knowledge, showing that the consequences of the dog attack caused an acute coronary thrombosis resulting in ST-segment elevation myocardial infarction in absence of bite-induced infection.

This study was partly supported by a grant of the Italian Ministry of Education, University and Research (MIUR): PON03PE_00009_4 'OPTIMA CARDIOPATHS'.